

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--	--

**Pearson Edexcel International Advanced Level**

**Wednesday 10 May 2023**

Morning (Time: 1 hour 30 minutes)

Paper  
reference

**WCH11/01**

**Chemistry**

**International Advanced Subsidiary/Advanced Level**

**UNIT 1: Structure, Bonding and Introduction to  
Organic Chemistry**

**You must have:**

Scientific calculator

Total Marks

### Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*

### Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

### Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P71860A

©2023 Pearson Education Ltd.  
N:1/1/1/1/



  
Pearson

## SECTION A

Answer ALL the questions in this section.

You should aim to spend no more than 20 minutes on this section.

For each question, select one answer from A to D and put a cross in the box . If you change your mind, put a line through the box  and then mark your new answer with a cross .

1 Which compound has the greatest covalent character?

- A MgBr<sub>2</sub>
- B MgF<sub>2</sub>
- C NaBr
- D NaF

(Total for Question 1 = 1 mark)

2 Which has the smallest ionic radius?

- A F<sup>-</sup>
- B Mg<sup>2+</sup>
- C Na<sup>+</sup>
- D O<sup>2-</sup>

(Total for Question 2 = 1 mark)

3 The first seven ionisation energies, in kJ mol<sup>-1</sup>, of an element are shown.

1010, 1900, 2910, 4960, 6270, 21 300, 25 400

In which group of the Periodic Table is this element located?

- A Group 3
- B Group 4
- C Group 5
- D Group 6

(Total for Question 3 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



4 What is the electronic configuration of an oxygen atom in its ground state?

	1s	2s	2p <sub>x</sub>	2p <sub>y</sub>	2p <sub>z</sub>	3s
<input type="checkbox"/> A	↑↓	↑↓	↑↓	↑↓		
<input type="checkbox"/> B	↑↓	↑↓	↑↑	↑	↑	
<input type="checkbox"/> C	↑↓	↑↓	↑	↑	↑	↑
<input type="checkbox"/> D	↑↓	↑↓	↑↓	↑	↑	

(Total for Question 4 = 1 mark)

5 Which property shows a **general decrease** across the Periodic Table from sodium to chlorine?

- A atomic radius
- B electronegativity
- C first ionisation energy
- D melting temperature

(Total for Question 5 = 1 mark)

6 The first ionisation energy of sulfur is lower than that of phosphorus.

Which is the best explanation for this?

- A the atomic radius of sulfur is greater than that of phosphorus
- B the electronegativity of sulfur is greater than that of phosphorus
- C the repulsion between the outer electrons of sulfur is greater than that of phosphorus
- D the shielding by the inner shell electrons of sulfur is greater than that of phosphorus

(Total for Question 6 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



7 1 kg of seawater contains 64 mg of bromide ions.

[ $A_r$  of Br = 80 Avogadro constant,  $L = 6.0 \times 10^{23} \text{ mol}^{-1}$ ]

(a) What is the concentration of bromide ions in parts per million (ppm) by mass? (1)


- A 0.80
- B 64
- C 800
- D 64 000

(b) How many bromide ions are in 500 g of the sample? (1)

- A  $2.4 \times 10^{20}$
- B  $4.8 \times 10^{20}$
- C  $1.9 \times 10^{22}$
- D  $3.8 \times 10^{22}$

(Total for Question 7 = 2 marks)

8 0.15 mol of aluminium is added to  $120 \text{ cm}^3$  of  $1.50 \text{ mol dm}^{-3}$  hydrochloric acid.



The amount of **unused** reactant is

- A 0.06 mol aluminium
- B 0.09 mol aluminium
- C 0.03 mol hydrochloric acid
- D 0.13 mol hydrochloric acid

(Total for Question 8 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



- 9 An organic compound contains 24.2% carbon, 4.00% hydrogen and 71.8% chlorine by mass.

[ $A_r$  values: H = 1.0 C = 12.0 Cl = 35.5]

What is the empirical formula of the compound?

- A CHCl
- B CH<sub>2</sub>Cl
- C C<sub>2</sub>H<sub>4</sub>Cl<sub>2</sub>
- D C<sub>4</sub>H<sub>4</sub>Cl<sub>4</sub>

(Total for Question 9 = 1 mark)

- 10 Which solution contains the **smallest** number of ions?

- A 500 cm<sup>3</sup> of 0.06 mol dm<sup>-3</sup> Ca(NO<sub>3</sub>)<sub>2</sub>(aq)
- B 500 cm<sup>3</sup> of 0.09 mol dm<sup>-3</sup> KI(aq)
- C 250 cm<sup>3</sup> of 0.12 mol dm<sup>-3</sup> BaCl<sub>2</sub>(aq)
- D 250 cm<sup>3</sup> of 0.09 mol dm<sup>-3</sup> (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>(aq)

(Total for Question 10 = 1 mark)

- 11 A piece of filter paper is soaked in water and attached to a microscope slide.

A few crystals of green copper(II) chromate(VI) are placed in the centre of the filter paper.

The filter paper is connected to a DC supply of 20V.

What colours are observed on the paper after a few minutes?

positive  
electrode



negative  
electrode

- A blue yellow
- B green blue
- C yellow green
- D yellow blue

(Total for Question 11 = 1 mark)

12 The melting temperature of beryllium is greater than that of barium.

What is the best explanation for this statement?

- A beryllium ions are smaller than barium ions
- B beryllium atoms have fewer outer shell electrons than barium atoms
- C beryllium ions have a smaller charge density than barium ions
- D beryllium atoms have a higher electronegativity than barium atoms

(Total for Question 12 = 1 mark)

13 Which molecule has the largest bond angle?

- A  $\text{BF}_3$
- B  $\text{CF}_4$
- C  $\text{H}_2\text{O}$
- D  $\text{NH}_3$

(Total for Question 13 = 1 mark)

14 Which molecule is polar?

- A  $\text{CO}_2$
- B  $\text{SF}_6$
- C  $\text{SO}_2$
- D  $\text{SiCl}_4$

(Total for Question 14 = 1 mark)

15 The reaction of methane with chlorine is a free radical substitution.


Which termination step does **not** occur?

- A  $\text{CH}_3\cdot + \text{CH}_3\cdot \rightarrow \text{C}_2\text{H}_6$
- B  $\text{CH}_3\cdot + \text{Cl}\cdot \rightarrow \text{CH}_3\text{Cl}$
- C  $\text{CH}_3\cdot + \text{H}\cdot \rightarrow \text{CH}_4$
- D  $\text{Cl}\cdot + \text{Cl}\cdot \rightarrow \text{Cl}_2$

(Total for Question 15 = 1 mark)



16 The repeat unit of a polymer is shown.




Which is the name of the monomer that forms this polymer?

- A 1,1,2-trimethylethene
- B 1,1-dimethylpropene
- C 2-methylbut-2-ene
- D 3-methylbut-2-ene

(Total for Question 16 = 1 mark)

17 What is the IUPAC name for the compound shown?




- A *E*-1-bromo-2-methylbut-1-ene
- B *Z*-1-bromo-2-methylbut-1-ene
- C *E*-1-bromo-2-ethyl-2-methylethene
- D *Z*-1-bromo-2-ethylpropene

(Total for Question 17 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



18 The skeletal formula of cyclohexa-1,4-diene is shown.



(a) The total number of  $\sigma$  (sigma) bonds in this molecule is

(1)

- A 4
- B 6
- C 12
- D 14

(b) 1.60 g of cyclohexa-1,4-diene is reduced to cyclohexane by reacting it with hydrogen gas.

[The molar volume of a gas is  $24.0 \text{ dm}^3 \text{ mol}^{-1}$  at room temperature and pressure (r.t.p.)

$M_r$  of cyclohexa-1,4-diene = 80]

What is the minimum volume, in  $\text{dm}^3$ , of hydrogen needed at r.t.p.?

(1)

- A 0.0400
- B 0.480
- C 0.960
- D 1.92

(Total for Question 18 = 2 marks)

TOTAL FOR SECTION A = 20 MARKS





DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

## SECTION B

Answer ALL the questions. Write your answers in the spaces provided.

19 This question is about mass spectrometry and the shapes of molecules.

(a) In a mass spectrometer vaporised atoms are ionised, and the ions formed are accelerated, deflected and detected.

(i) State how atoms are ionised in the mass spectrometer.

(1)

.....

.....

.....

(ii) State how the ions formed are accelerated.

(1)

.....

.....

(iii) Explain why isotopes of an element have the same chemical reactions but their ions are deflected differently in a mass spectrometer.

(2)

.....

.....

.....

.....

.....

(b) Data from mass spectra may be used to determine the relative atomic masses of elements.

(i) State what is meant by relative atomic mass.

(2)

.....

.....

.....

.....




(ii) A sample of chlorine contains 75.53 % of  $^{35}\text{Cl}$  and 24.47 % of  $^{37}\text{Cl}$ .

Calculate the relative atomic mass of this sample of chlorine, giving your answer to **four** significant figures.

(2)

(c) The mass spectrum of phosphorus trichloride,  $\text{PCl}_3$ , is shown.

Phosphorus has only one isotope,  $^{31}\text{P}$ .



(i) There are three peaks in the region of  $m/z = 101-105$ .

Complete the table to show the ions responsible for these peaks.

(2)

$m/z$	Formula of ion	Relative peak height
101		9
103		6
105		1



(ii) Show that the relative peak heights given in the table are consistent with the isotopic ratio of  $^{35}\text{Cl}$  to  $^{37}\text{Cl}$  being 3:1.

(2)

.....

.....

.....

(d) (i) Draw a dot-and-cross diagram of a  $\text{PCl}_3$  molecule.  
Show outer electrons only.

(2)

(ii) Explain the shape of a  $\text{PCl}_3$  molecule.

(3)

.....

.....

.....

.....

.....

.....

.....

**(Total for Question 19 = 17 marks)**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 7 1 8 6 0 A 0 1 1 2 4

- 20 (a) Barium chloride,  $\text{BaCl}_2$ , can be prepared by the reaction of barium carbonate with hydrochloric acid.



- (i) Write the ionic equation for this reaction.  
State symbols are not required.

(1)

- (ii) Calculate the atom economy by mass for this preparation of barium chloride.

[ $A_r$  values: H = 1.0 C = 12.0 O = 16.0 Cl = 35.5 Ba = 137.3]

(2)



- (iii) Barium chloride has a melting temperature of 962 °C.  
Caesium chloride has a melting temperature of 646 °C.

Explain, by considering the ions involved, the difference between the melting temperatures of these Period 6 chlorides.

(4)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (iv) Covalent and ionic bonding are the extremes of a continuum of bonding type.

Explain the difference in bonding between barium chloride and beryllium chloride, using the electronegativity values shown.

Element	Electronegativity
Ba	0.9
Be	1.5
Cl	3.0

(2)

.....

.....

.....

.....


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) Barium chloride reacts with Group 1 sulfates,  $M_2SO_4$ .



A solution is made by dissolving 7.98 g of a Group 1 sulfate in deionised water.

Excess aqueous barium chloride is added to this solution and the precipitate is filtered, dried and weighed.

The mass of the barium sulfate precipitate is 10.72 g.

Identify the Group 1 element.

[ $A_r$  values: Ba = 137.3    S = 32.1    O = 16.0]


(4)

(Total for Question 20 = 13 marks)



21 Graphite electrodes are used in the extraction of aluminium by the electrolysis of aluminium oxide,  $\text{Al}_2\text{O}_3$ , dissolved in a solvent (molten cryolite) at a temperature of  $950^\circ\text{C}$ .

- (a) Explain how the structure and bonding in graphite make it suitable for this application.  
Refer to the diagram in your answer.



graphite

(3)

.....

.....

.....

.....

.....

.....


.....



(b) The half-equations for the electrolysis reactions are shown.



The oxygen produced reacts with the graphite electrode.



- (i) Calculate the maximum volume, in  $\text{dm}^3$ , of  $\text{CO}_2$ , measured at r.t.p. which could be produced when 1.00 kg aluminium is extracted using this process.

[The molar volume of a gas is  $24.0 \text{ dm}^3 \text{ mol}^{-1}$  at r.t.p.]

(4)





(ii) Every year in the United Kingdom, about 7.2 billion aluminium cans are recycled.

Suggest three ways this recycling reduces energy consumption.

(3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**(Total for Question 21 = 10 marks)**

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



**22** Crude oil is mainly a mixture of saturated hydrocarbons that can be separated by fractional distillation.

The diagram shows, in a simplified form, the products of fractional distillation.



(a) (i) State what is meant by a saturated hydrocarbon.

(2)

.....

.....

.....

.....

(ii) Describe how the fractions in crude oil are separated during fractional distillation.

(2)

.....

.....

.....

.....



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

- (b) Butane,  $C_4H_{10}$ , is found in the refinery gases fraction of crude oil.  
It is used as a propellant in pharmaceutical inhalers.  
An inhaler that provides a total of 120 doses contains 1.55 g of butane.

Calculate the volume, in  $cm^3$ , of propellant used for each dose  
at  $25^\circ C$  and 100 kPa.

Use the ideal gas equation and give your answer to an appropriate number of  
significant figures.

$$[pV = nRT \quad R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}]$$

(5)



(c) Octane,  $C_8H_{18}$ , is found in the gasoline fraction of crude oil.

- (i) Write an equation for the complete combustion of octane.  
Include state symbols.

(2)

- (ii) Give **two** reasons why alternative fuels, such as bioethanol, are being developed to replace those produced from crude oil.

(2)

.....

.....

.....

.....

.....

.....

- (iii) Straight-chain alkanes, such as octane, are converted into branched-chain alkanes to improve the performance of petrol in car engines.

Name this conversion process.

(1)

- (iv) Give an equation using skeletal formulae for the conversion of octane into 2,5-dimethylhexane.  
State symbols are not required.

(1)



(d) In steam cracking, a mixture of hydrocarbons and steam is heated to 850°C for a very short time in the absence of oxygen.

(i) Give a reason why oxygen should be excluded during steam cracking.

(1)

(ii) Ethene is the major product.

State one use of ethene.


(1)

(e) Ethene reacts with hydrogen bromide to form bromoethane.

Draw the mechanism for this reaction.

Include curly arrows, and relevant lone pairs and dipoles.

(3)



(Total for Question 22 = 20 marks)

**TOTAL FOR SECTION B = 60 MARKS**  
**TOTAL FOR PAPER = 80 MARKS**



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**BLANK PAGE**



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**BLANK PAGE**



# The Periodic Table of Elements

1 2 3 4 5 6 7 0 (8) (18)

1.0	<b>H</b>	hydrogen	1
-----	----------	----------	---

**Key**

relative atomic mass
<b>atomic symbol</b>
name
atomic (proton) number

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
6.9	9.0	45.0	47.9	50.9	52.0	54.9	55.8	58.9	58.7	63.5	65.4	10.8	12.0	14.0	16.0	19.0	4.0
<b>Li</b>	<b>Be</b>	<b>Sc</b>	<b>Ti</b>	<b>V</b>	<b>Cr</b>	<b>Mn</b>	<b>Fe</b>	<b>Co</b>	<b>Ni</b>	<b>Cu</b>	<b>Zn</b>	<b>B</b>	<b>C</b>	<b>N</b>	<b>O</b>	<b>F</b>	<b>He</b>
lithium	beryllium	scandium	titanium	vanadium	chromium	manganese	iron	cobalt	nickel	copper	zinc	boron	carbon	nitrogen	oxygen	fluorine	helium
3	4	21	22	23	24	25	26	27	28	29	30	5	6	7	8	9	2
23.0	24.3	88.9	91.2	92.9	95.9	[98]	101.1	102.9	106.4	107.9	112.4	27.0	28.1	31.0	32.1	35.5	39.9
<b>Na</b>	<b>Mg</b>	<b>Y</b>	<b>Zr</b>	<b>Nb</b>	<b>Mo</b>	<b>Tc</b>	<b>Ru</b>	<b>Rh</b>	<b>Pd</b>	<b>Ag</b>	<b>Cd</b>	<b>Al</b>	<b>Si</b>	<b>P</b>	<b>S</b>	<b>Cl</b>	<b>Ar</b>
sodium	magnesium	yttrium	zirconium	niobium	molybdenum	technetium	ruthenium	rhodium	palladium	silver	cadmium	aluminium	silicon	phosphorus	sulfur	chlorine	argon
11	12	39	40	41	42	43	44	45	46	47	48	13	14	15	16	17	18
39.1	40.1	88.9	91.2	92.9	95.9	[98]	101.1	102.9	106.4	107.9	112.4	69.7	72.6	74.9	79.0	79.9	83.8
<b>K</b>	<b>Ca</b>	<b>La*</b>	<b>Hf</b>	<b>Ta</b>	<b>W</b>	<b>Re</b>	<b>Os</b>	<b>Ir</b>	<b>Pt</b>	<b>Au</b>	<b>Hg</b>	<b>Ga</b>	<b>Ge</b>	<b>As</b>	<b>Se</b>	<b>Br</b>	<b>Kr</b>
potassium	calcium	lanthanum	hafnium	tantalum	tungsten	rhenium	osmium	iridium	platinum	gold	mercury	gallium	germanium	arsenic	selenium	bromine	krypton
19	20	57	72	73	74	75	76	77	78	79	80	31	32	33	34	35	36
85.5	87.6	138.9	178.5	180.9	183.8	186.2	190.2	192.2	195.1	197.0	200.6	69.7	72.6	74.9	79.0	79.9	131.3
<b>Rb</b>	<b>Sr</b>	<b>La*</b>	<b>Hf</b>	<b>Ta</b>	<b>W</b>	<b>Re</b>	<b>Os</b>	<b>Ir</b>	<b>Pt</b>	<b>Au</b>	<b>Hg</b>	<b>In</b>	<b>Sn</b>	<b>Sb</b>	<b>Te</b>	<b>I</b>	<b>Xe</b>
rubidium	strontium	lanthanum	hafnium	tantalum	tungsten	rhenium	osmium	iridium	platinum	gold	mercury	indium	tin	antimony	tellurium	iodine	xenon
37	38	57	72	73	74	75	76	77	78	79	80	49	50	51	52	53	54
132.9	137.3	138.9	178.5	180.9	183.8	186.2	190.2	192.2	195.1	197.0	200.6	114.8	118.7	121.8	127.6	126.9	131.3
<b>Cs</b>	<b>Ba</b>	<b>La*</b>	<b>Hf</b>	<b>Ta</b>	<b>W</b>	<b>Re</b>	<b>Os</b>	<b>Ir</b>	<b>Pt</b>	<b>Au</b>	<b>Hg</b>	<b>Pb</b>	<b>Bi</b>	<b>Po</b>	<b>At</b>	<b>Rn</b>	<b>Rn</b>
caesium	barium	lanthanum	hafnium	tantalum	tungsten	rhenium	osmium	iridium	platinum	gold	mercury	lead	bismuth	polonium	astatine	radon	radon
55	56	57	72	73	74	75	76	77	78	79	80	82	83	84	85	86	86
[223]	[226]	[227]	[261]	[262]	[266]	[264]	[277]	[268]	[271]	[272]	[272]	204.4	207.2	209.0	[210]	[222]	[222]
<b>Fr</b>	<b>Ra</b>	<b>Ac*</b>	<b>Rf</b>	<b>Db</b>	<b>Sg</b>	<b>Bh</b>	<b>Hs</b>	<b>Mt</b>	<b>Ds</b>	<b>Rg</b>	<b>Rg</b>	<b>Tl</b>	<b>Pb</b>	<b>Bi</b>	<b>Po</b>	<b>At</b>	<b>Rn</b>
francium	radium	actinium	rutherfordium	dubnium	seaborgium	bohrium	hassium	meitnerium	darmstadtium	roentgenium	roentgenium	thallium	lead	bismuth	polonium	astatine	radon
87	88	89	104	105	106	107	108	109	110	111	111	81	82	83	84	85	86

Elements with atomic numbers 112-116 have been reported but not fully authenticated

140	141	144	150	152	157	163	165	167	169	173	175
<b>Ce</b>	<b>Pr</b>	<b>Nd</b>	<b>Sm</b>	<b>Eu</b>	<b>Gd</b>	<b>Dy</b>	<b>Ho</b>	<b>Er</b>	<b>Tm</b>	<b>Yb</b>	<b>Lu</b>
cerium	praseodymium	neodymium	samarium	europium	gadolinium	dysprosium	holmium	erbium	thulium	ytterbium	lutetium
58	59	60	62	63	64	66	67	68	69	70	71
232	[231]	238	[242]	[243]	[247]	[251]	[254]	[253]	[256]	[254]	[257]
<b>Th</b>	<b>Pa</b>	<b>U</b>	<b>Pu</b>	<b>Am</b>	<b>Cm</b>	<b>Cf</b>	<b>Es</b>	<b>Fm</b>	<b>Md</b>	<b>No</b>	<b>Lr</b>
thorium	protactinium	uranium	plutonium	americium	curium	californium	einsteinium	fermium	mendeleevium	nobelium	lawrencium
90	91	92	94	95	96	98	99	100	101	102	103

\* Lanthanide series

\* Actinide series

