



Mark Scheme (Results)

October 2023

Pearson Edexcel International Advanced
Subsidiary Level In Chemistry (WCH13)
Paper 01
Unit 3: Practical Skills in Chemistry I

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

October 2023

Question Paper Log Number: P75062A

Publications Code: WCH13_01_MS_2310

All the material in this publication is copyright.

© Pearson Education Ltd 2023

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.


Question Number	Answer	Additional Guidance	Mark
1(a)(i)	<p>A description that makes reference to the following points:</p> <ul style="list-style-type: none"> • use of nichrome / platinum / Pt wire • use of (concentrated) hydrochloric acid / HCl(aq) • flame test method 	<p>(1) Allow nickel-chromium / Ni-Cr Allow silica rod Allow rod for wire Do not award just 'nickel' / just 'chromium' Do not award wooden splint</p> <p>(1) Allow any mention of HCl(aq) e.g. cleaning or mixing solid and acid or making a paste / solution Allow HCl for HCl(aq) Ignore dilute Do not award other acids</p> <p>(1) (wire then) dipped in solid and (placed) in (hot / roaring / colourless / blue-cone / non-luminous) (Bunsen) flame</p> <p>Allow salt / compound / substance / paste / sample / solution for 'solid' Allow on / over / under / near / show / above for 'in' Allow spirit / ethanol burner</p> <p>Do not award 'metal' for solid Do not award fire for flame Do not award yellow / luminous flame Ignore 'burn'</p>	(3)

Question Number	Answer	Additional Guidance	Mark
1(a)(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> statement of both flame colours identification of Ba²⁺ 	<p>(1) Na⁺ = (persistent) yellow Allow gold / orange / yellow-orange K⁺ = lilac Allow (pale) purple Do not award violet</p> <p>(1) Ignore barium / barium ion Do not award Cu²⁺</p>	(2)

Question Number	Answer	Additional Guidance	Mark
1(b)(i)	<p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> chloride / Cl⁻ and bromide / Br⁻ and iodide / I⁻ 	<p>Allow omission of the charge once Ignore chlorine ion, bromine ion or iodine ion Do not award just chlorine, bromine or iodine</p>	(1)

Question Number	Answer	Additional Guidance	Mark
1(b)(ii)	<p>An answer that makes reference to the following points:</p> <p>EITHER</p> <ul style="list-style-type: none"> • addition of dilute (aqueous) ammonia (1) • only silver chloride / chloride (precipitate) dissolves (1) • addition of concentrated (aqueous) ammonia (1) • only silver iodide / iodide is insoluble (1) <p>OR</p> <ul style="list-style-type: none"> • addition of concentrated sulfuric acid /H₂SO₄ (1) • silver chloride / chloride precipitate gives steamy / misty / white fumes only (1) • silver bromide / bromide precipitate gives brown fumes (1) • silver iodide / iodide precipitate gives purple vapour fumes / gas (1) 	<p>Penalise use of halogen for halide once only Allow use of halogen ion / anion The sequence suggested must allow the ions to be distinguished</p> <p>Allow bromide and iodide do not dissolve</p> <p>Accept only silver bromide / bromide precipitate dissolves if chloride has been eliminated and conc ammonia added to remaining precipitates</p> <p>If no other mark is scored ‘addition of (aqueous) ammonia scores (1)</p> <p>Allow addition to A, B and C for this set of tests</p> <p>Allow choking gas evolved / choking smell</p> <p>Allow black / grey solid or ‘bad eggs’ smell Allow yellow solid (of sulfur) Ignore choking gas evolved / choking smell</p> <p>If no other mark is scored, silver chloride is white, and silver bromide is cream and silver iodide is yellow scores (1)</p>	(4)

(Total for Question 1 = 10 marks)

Question Number	Answer	Additional Guidance	Mark
2(a)	<p>Diagram showing the following points</p> <ul style="list-style-type: none"> • boiling tube • delivery tube above the level of the liquid in the boiling tube if shown or above half the height of the tube • sealed connection between reaction vessel and delivery tube • inverted measuring cylinder containing some water (shown as a line or an annotation indicating it is full) • water level in trough over the bottom of the measuring cylinder • end of delivery tube below or in the measuring cylinder <p>6 points scores 3 marks; 5 or 4 points scores 2 marks; 2 or 3 points scores 1 mark</p>	<p>Allow any reaction apparatus that can be fitted with a bung and delivery tube Allow any part of a delivery tube shown</p> <p>Allow bung not shown as cross-section</p> <p>Allow measuring cylinder shown not vertical Allow other collecting vessels with graduations</p> <p>Ignore omission of graduations on measuring cylinder Ignore omission of beehive shelf Ignore gas syringe</p> <p>Ignore the delivery tube passing through the wall of the trough</p> <p style="text-align: center;">Alternative allowed bungs</p>  <p>The diagram shows a boiling tube containing blue liquid, with a delivery tube extending from its mouth down into an inverted measuring cylinder. The measuring cylinder is partially filled with blue liquid and is submerged in a trough of water. To the right, two alternative bung designs are shown: one is a simple orange square with two vertical lines, and the other is an orange square with two vertical lines and a central vertical slot.</p> <p>Ignore apparatus labels including 'heat'</p>	(3)

Question Number	Answer	Additional Guidance	Mark
2(b)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> ethanoic acid is a weak acid (and hydrochloric acid is a strong acid) (1) (with hydrochloric acid) <p>gas would escape before the boiling tube was sealed (1)</p>	<p>Allow 'hydrochloric acid is a stronger acid'</p> <p>Allow reaction would be slow</p> <p>Allow reaction would be less vigorous</p> <p>Allow reaction would not be violent</p> <p>Allow reverse arguments</p> <p>Ignore just 'hydrochloric acid is a strong acid'</p> <p>Allow just 'less gas would escape'</p>	(2)

Question Number	Answer	Additional Guidance	Mark
2(c)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> to take account of any calcium carbonate left in the weighing bottle 	<p>Ignore just 'gives the mass of CaCO₃ that reacts'</p> <p>Ignore just 'gives exact amount of CaCO₃'</p>	(1)

Question Number	Answer	Additional Guidance	Mark
2(d)(i)	<ul style="list-style-type: none"> • correct axes with at least 50% of the grid used in both directions (1) • axes labelled with correct units (1) • all points plotted correctly (1) 	<p>mass on x axis; volume on y axis Allow scale 0.10 g = 1 big square</p> <p>Allow grammatical errors e.g. use of brackets rather than ‘/’ before units Allow mass / g and vol / cm³ for labels</p> <p>Allow plotting to within half a small square</p> <p>TE for M2 and M3 if axes wrong way round</p>	<p>(3)</p> <p>Clip with d(ii) and (d)(iii)</p>

Question Number	Answer	Additional Guidance	Mark
2(d)(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> • best fit line drawn through five points and passes through the origin • structure line from 0.25 g to graph and line to 52.5 cm³ 	<p>(1) Do not award BFL drawn to include value at 0.2 g Allow TE on points plotted in (d)(i) even if it does not go through origin Accept line that stops at 0.11 g data point but would pass through the origin if extrapolated Ignore extrapolation beyond 0.36 / 76</p> <p>(1) Allow vol at 0.25 g marked on BFL Allow 51.5-53.5 cm³ TE only on any straight line covering all the points</p> <p>Ignore calculation of volume</p>	<p>(2)</p> <p>Clip with d(i) and d(iii)</p>

Question Number	Answer	Additional Guidance	Mark
2(d)(iii)	<ul style="list-style-type: none"> • calculation of molar mass of CaCO₃ (1) • calculation of amount of CaCO₃ (1) • scale volume of CO₂ to 1 mol (1) 	<p>Example of calculation</p> <p>molar mass = $40.1 + 12 + 16 \times 3 = 100.1 \text{ (g mol}^{-1}\text{)}$</p> <p>mol (CaCO₃) = $0.25 \div 100.1 = 2.4975 \times 10^{-3} / 0.0024975$</p> <p>1 mol CO₂ occupies $52.5 \div 2.4975 \times 10^{-3} = 21021 \text{ (cm}^3\text{)} / 21.021 \text{ dm}^3$ Units must be correct if given but accept $\text{cm}^3 \text{ mol}^{-1} / \text{dm}^3 \text{ mol}^{-1}$</p> <p>TE at each stage and on volume in (d)(ii)</p> <p>Ignore SF except 1 SF</p> <p>Correct answer with some working scores (3)</p> <p>Allow use of $M_r \text{ (CaCO}_3\text{)} = 100 \text{ (} V_m = 21\text{)}$</p> <p>Use of 51.5 gives 20621 cm³ (20600 for $M_r = 100$) Use of 52 gives 20821 cm³ (20800 for $M_r = 100$) Use of 53 gives 21221 cm³ (21200 for $M_r = 100$) Use of 53.5 gives 21421 cm³ (21400 for $M_r = 100$)</p> <p>Calculation of moles of acid (0.030) divided by 2 (0.015) Vol of CO₂ $\div 0.015$ e.g. $52.5 \div 0.015 = 3500 \text{ cm}^3$ scores M3 only</p>	<p>(3)</p> <p>Clip with d(i) and d(ii)</p>

Question Number	Answer	Additional Guidance	Mark
2(e)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> • some carbon dioxide / CO₂ / gas will escape before the boiling tube can be sealed • some carbon dioxide / gas will dissolve in the water 	<p>(1) Allow Some CO₂ / gas will escape in Step 5 Allow Some CO₂ / gas will escape when the solid is added Allow reaction starts before the boiling tube can be sealed</p> <p>Ignore just 'Some carbon dioxide / gas will escape'</p> <p>(1) Accept carbon dioxide is soluble (in water)</p> <p>Ignore reference to temperature or pressure not rtp Ignore 'some CO₂ remains in apparatus' Ignore 'incomplete reaction'</p> <p>Do not award references to measurement errors loss of reactant apparatus damaged or not working properly</p>	(2)


(Total for Question 2 = 16 marks)

Question Number	Answer	Additional Guidance	Mark
3(a)(i)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> flammable symbol identified <p>and</p> <ul style="list-style-type: none"> harmful to the environment symbol identified 	<p>Allow inflammable Ignore combusts / burns easily</p> <p>Allow alternatives to 'harmful to' e.g. 'damages' / 'bad for' / 'poisonous' / 'toxic' / 'hazard' to the environment Allow 'living things' / 'organisms' for 'the environment' Ignore type of environment e.g. aquatic Ignore pollutes the environment</p> <p>Do not award 'biohazard' Do not award symbols the wrong way round Do not award just 'toxic' / 'poisonous' / 'harmful'</p>	<p>(1)</p> <p>Clip with (a)(ii)</p>

Question Number	Answer	Additional Guidance	Mark
3(a)(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> (cyclohexanol(flammable)) use an electric heater (1) (cyclohexene (harmful to the environment)) use an organic waste bottle / separate container (1) 	<p>Ignore use of fume cupboard, goggles, lab coat</p> <p>Accept isomantle Allow water bath / oil bath Allow 'no (naked) flame / fire' Ignore 'keep away from oxidising agents' Ignore keep away from Bunsen burner</p> <p>Accept do not pour the (organic) waste down the sink / drain Allow do not release into the environment</p> <p>Allow one use of 'use small amounts' in M1 or M2 No TE on incorrect identification of hazard symbols</p>	<p>(2)</p>

Question Number	Answer	Additional Guidance	Mark
3(b)	<p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> (anti-bumping granules) provides a nucleus on which gas bubbles grow 	<p>Accept provide a surface /site for bubbles to form on Allow prevent local heating / superheating Allow 'distribute the heat' Allow prevent the (sudden) production of large gas bubbles (which cause bumping) Allow ensures that gas bubbles are small Ignore stir the reaction mixture Ignore prevent flash boiling / sudden boiling</p>	(1)

Question Number	Answer	Additional Guidance	Mark
3(c)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> (fractional distillation) gives better separation of the substances in the reaction flask (than simple distillation) (fractional distillation column) gives longer reaction time <p>OR</p> <ul style="list-style-type: none"> 'better separation of cyclohexene and water' scores 2 marks 	<p>(1) Allow (fractional distillation) is more effective / efficient (than simple distillation) Allow just 'better separation' is achieved Allow purer product obtained Ignore reference to increased yield Ignore 'more accurate' Ignore 'similar boiling temperatures'</p> <p>(1) Accept fractional distillation column acts like a reflux condenser Accept cyclohexanol has a higher boiling temperature (than cyclohexene) so is returned to the flask, increasing reaction time / yield Allow to give (more) complete reaction</p> <p>Just 'separation of cyclohexene and water' scores (1)</p>	(2)

Question Number	Answer	Additional Guidance	Mark
3(d)	<p>A diagram that shows the following:</p> <ul style="list-style-type: none"> separating funnel top layer labelled cyclohexene / organic and lower layer labelled water / aqueous 	<p>(1) The funnel must have a tap and have a neck capable of being closed with a stopper or a bung Allow round separating funnel Allow bung / stopper not shown Ignore tap at the top</p> <p>Do not award a filter funnel Do not award funnel without some sort of tapering at the neck</p> <p>Allow for (2)</p>  <p>(1)</p> <p>Allow minor error in organic name e.g. cyclohexane Ignore 'inorganic lower layer' Do not award if cyclohexanol is in the lower layer</p>	(2)

Question Number	Answer	Additional Guidance	Mark
3(e)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> (the organic layer) changes from cloudy to clear (anhydrous calcium chloride) removes the (traces of) water 	<p>(1) Allow just 'becomes clear / transparent' Ignore 'less cloudy', clearer, more transparent Ignore white Ignore colourless Ignore the drying agent clumps together Ignore layers disappear</p> <p>(1) Accept (anhydrous calcium chloride) dries the cyclohexene Allow (anhydrous) calcium chloride is a drying agent Ignore calcium chloride becomes hydrated</p>	(2)

Question Number	Answer	Additional Guidance	Mark
3(f)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> lower temperature within a suitable range and upper temperature within a suitable range 	<p>79–82 (°C)</p> <p>84–88 (°C)</p> <p>Do not award a range which starts or ends with 83 (°C)</p>	(1)

Question Number	Answer	Additional Guidance	Mark
3 (g)	<ul style="list-style-type: none"> • calculation of mass of cyclohexene formed from 3.96 g of cyclohexanol with 100% yield (1) • calculation of percentage yield (1) 	<p>Example of calculation</p> <p>100 g of cyclohexanol forms 82 g of cyclohexene and (so) 3.96 g forms $3.96 \times 82 \div 100 = 3.2472$ (g)</p> <p>% yield = $100 \times 2.09 \div 3.2472 = 64.363\%$</p> <p>Method using calculation of moles mol of cyclohexanol = $3.96 \div 100$ = $3.96 \times 10^{-2} / 0.0396$ = mol cyclohexene mass of cyclohexene for 100% yield = $82 \times 0.0396 = 3.2472$ (g) (1) % yield = $100 \times 2.09 \div 3.2472$ = 64.363% (1)</p> <p>Ignore SF except 1 SF</p> <p>TE for numerical errors in M1 unless %>100</p> <p>Allow any correct method Correct answer with some working scores (2)</p> <p>$100 \times 2.09 \div 3.96 = 52.778\%$ scores zero</p>	(2)

Question Number	Answer	Additional Guidance	Mark											
3(h)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> • any three correct observations (1) • fourth correct observation (1) 	<table border="1" data-bbox="1189 304 1924 799"> <thead> <tr> <th data-bbox="1189 304 1413 376" rowspan="2">Test</th> <th colspan="2" data-bbox="1413 304 1924 376">Observations</th> </tr> <tr> <th data-bbox="1413 376 1666 443">cyclohexanol</th> <th data-bbox="1666 376 1924 443">cyclohexene</th> </tr> </thead> <tbody> <tr> <td data-bbox="1189 443 1413 624">addition of phosphorus(V) chloride</td> <td data-bbox="1413 443 1666 624">steamy / misty / white fumes</td> <td data-bbox="1666 443 1924 624">no change / no reaction / no observation</td> </tr> <tr> <td data-bbox="1189 624 1413 799">addition of bromine water</td> <td data-bbox="1413 624 1666 799">brown / orange / yellow Br₂(aq) unchanged</td> <td data-bbox="1666 624 1924 799">brown / orange / yellow Br₂(aq) turns colourless</td> </tr> </tbody> </table> <p data-bbox="1178 839 1637 871">For PCl₅ do not award white smoke</p> <p data-bbox="1178 916 1834 979">For Br₂(aq) and cyclohexanol allow no change / no reaction / no observation</p> <p data-bbox="1178 1024 1877 1088">For Br₂(aq) and cyclohexene allow just 'decolorised' / 'turns colourless'</p> <p data-bbox="1178 1101 1843 1133">Do not award red or red-brown for colour of Br₂(aq)</p> <p data-bbox="1178 1171 1733 1203">Do not award additional incorrect observations</p>	Test	Observations		cyclohexanol	cyclohexene	addition of phosphorus(V) chloride	steamy / misty / white fumes	no change / no reaction / no observation	addition of bromine water	brown / orange / yellow Br ₂ (aq) unchanged	brown / orange / yellow Br ₂ (aq) turns colourless	(2)
Test	Observations													
	cyclohexanol	cyclohexene												
addition of phosphorus(V) chloride	steamy / misty / white fumes	no change / no reaction / no observation												
addition of bromine water	brown / orange / yellow Br ₂ (aq) unchanged	brown / orange / yellow Br ₂ (aq) turns colourless												

(Total for Question 3 = 15 marks)

Question Number	Answer	Additional Guidance	Mark
4(a)	An answer that makes reference to the following points: <ul style="list-style-type: none"> <li data-bbox="371 347 1196 384">• (from) yellow (1) <li data-bbox="371 421 1196 458">• (to) orange (1) 	Do not award 'red / pink' From orange to yellow scores (1)	(2)

Question Number	Answer	Additional Guidance	Mark
4(b)	An answer that makes reference to the following: <ul style="list-style-type: none"> <li data-bbox="371 756 573 793">• 23.40 (cm³) 	Allow 23.4 (cm ³)	(1)

Question Number	Answer	Additional Guidance	Mark
4(c)	<p>A description that makes reference to the following points:</p> <ul style="list-style-type: none"> • add the acid (quickly) until just short of the rough end-point • add the acid drop-by-drop • with swirling <p>and</p> <p>until the indicator colour (just) changes</p>	<p>(1) Accept within 1– 4 cm³ (any value in this range) before the rough titre Allow to a value in the range 19.4 – 22.4 cm³ Allow ‘until close to the rough value’ Ignore ‘carbonate added’ Do not award ‘until the rough value is reached’</p> <p>(1) Standalone mark (award even if M1 not given) Allow ‘dropwise’ Allow ‘a few drops at a time’ Ignore ‘add very slowly’</p> <p>Allow any indication of mixing Allow shaking Allow stirring</p> <p>(1) Allow until the end-point Allow any stated final colour</p> <p>Ignore references to filling the burette, use of the pipette, white tiles</p>	<p>(3)</p> <p>Clip with (b)</p>

Question Number	Answer	Additional Guidance	Mark
4(d)	<ul style="list-style-type: none"> • calculation of moles of sodium carbonate (1) • use of 2:1 ratio to gives moles of HCl in 22.65 cm³ (1) • concentration of hydrochloric acid in mol dm⁻³ (1) 	<p>Example of calculation</p> $\text{mol Na}_2\text{CO}_3 = 25.0 \times 0.105 \div 1000$ $= 2.625 \times 10^{-3} / 0.002625$ $2 \times 2.625 \times 10^{-3} = 5.25 \times 10^{-3}$ $5.25 \times 10^{-3} \times 1000 \div 22.65$ $= 0.23179 / 2.3179 \times 10^{-1} \text{ (mol dm}^{-3}\text{)}$ <p>The reacting volumes transposed gives the final concentration of hydrochloric acid = 0.19026 (mol dm⁻³). This scores (2)</p> <p>The same volume used twice will give the final concentration of hydrochloric acid = 0.2100 (mol dm⁻³). These score (2)</p> <p>TE at each stage Ignore SF except 1 SF Allow any correct method Correct answer with some working scores (3)</p> <p>Comment If working shows factor of 1000 omitted twice in the step by step calculation max (2)</p>	(3)

(Total for Question 4 = 9 marks)
(Total for Paper = 50 marks)

