



Mark Scheme (Results)

October 2024

Pearson Edexcel International Advanced Level
In Chemistry (WCH16)
Paper 01 Practical Skills in Chemistry II

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

October 2024

Question Paper Log Number P78394A

Publications Code WCH16_01_2410_MS

All the material in this publication is copyright

© Pearson Education Ltd 2024

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Using the Mark Scheme

Examiners should look for qualities to reward rather than faults to penalise. This does NOT mean giving credit for incorrect or inadequate answers, but it does mean allowing candidates to be rewarded for answers showing correct application of principles and knowledge. Examiners should therefore read carefully and consider every response: even if it is not what is expected it may be worthy of credit.

The mark scheme gives examiners:

- an idea of the types of response expected
- how individual marks are to be awarded
- the total mark for each question
- examples of responses that should NOT receive credit.

/ means that the responses are alternatives and either answer should receive full credit.

() means that a phrase/word is not essential for the award of the mark, but helps the examiner to get the sense of the expected answer.

Phrases/words in **bold** indicate that the meaning of the phrase or the actual word is **essential** to the answer.

ecf/TE/cq (error carried forward) means that a wrong answer given in an earlier part of a question is used correctly in answer to a later part of the same question.

Candidates must make their meaning clear to the examiner to gain the mark. Make sure that the answer makes sense. Do not give credit for correct words/phrases which are put together in a meaningless manner. Answers must be in the correct context.

Quality of Written Communication

Questions which involve the writing of continuous prose will expect candidates to:

- write legibly, with accurate use of spelling, grammar and punctuation in order to make the meaning clear
- select and use a form and style of writing appropriate to purpose and to complex subject matter
- organise information clearly and coherently, using specialist vocabulary when appropriate.

Full marks will be awarded if the candidate has demonstrated the above abilities. Questions where QWC is likely to be particularly important are indicated (QWC) in the mark scheme, but this does not preclude others.

Question Number	Answer	Additional Guidance	Mark
1(a)(i)	An answer that makes reference to the following points: <ul style="list-style-type: none"> • AgI/silver iodide • NH₃/ammonia • Fe(OH)₃/iron(III)hydroxide/ Fe(OH)₃·(H₂O)₃ • BaSO₄/barium sulfate 	<p>If both name and formula are given, both must be correct</p> <p>(1)</p> <p>(1) Do not award ammonium/NH₄⁺</p> <p>(1) Do not award Fe(OH)₂/iron(II)hydroxide</p> <p>(1) Allow sulphate</p>	(4)

Question Number	Answer	Additional Guidance	Mark
1(a)(ii)	<ul style="list-style-type: none"> • barium iodide / BaI₂ 	<p>If name and formula are given, both must be correct</p> <p>Ignore any state symbols Do not award just Ba²⁺/ barium</p>	(1)

Question Number	Answer	Additional Guidance	Mark
1(a)(iii)	An explanation that makes reference to the following points: <ul style="list-style-type: none"> • Fe(III)/Fe³⁺ and NH₄⁺ and SO₄²⁻ • Fe NH₄ (SO₄)₂ 	<p>(1)</p> <p>(1) Accept cations in either order Allow formula with water of crystallisation. Ignore state symbols Ignore incorrect brackets e.g.(NH₄) TE from incorrect ions in M1</p>	(2)

Question Number	Answer	Additional Guidance	Mark
1(b)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> iodine has been produced (from iodide ions) Fe(III) is reduced /Iron is reduced from +3 to +2 <p>or</p> <p>I⁻ oxidised/iodine is oxidised from -1 to 0</p>	<p>(1) Allow I⁻ changes to I₂</p> <p>(1) Iodide ions oxidised to iodine scores both marks Allow Fe(III) acts as an oxidising agent/I⁻ acts as reducing agent</p> <p>Ignore references to the identity of the white ppt Do not award unbalanced (half)equations for M2</p>	(2)

(Total for Question 1 = 9 marks)

Question Number	Answer	Additional Guidance	Mark
2(a)(i)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> burette or graduated pipette or pipettes with markings 	<p>Do not award just pipette Do not award volumetric pipette/flask Do not award measuring cylinder</p>	(1)

Question Number	Answer	Additional Guidance	Mark
2(a)(ii)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> blue/black/blue-black/dark blue 	<p>Allow colourless to blue/black Allow black-blue Do not award colourless/yellow/brown/purple</p>	(1)

Question Number	Answer	Additional Guidance	Mark
2(a)(iii)	An answer that makes reference to the following point: <ul style="list-style-type: none"> the rate of reaction does not change significantly/is constant at the start of the reaction 	<p>Allow (so that)any change in concentration of the peroxydisulfate/iodide does not affect the reaction rate</p> <p>Allow so that the thiosulfate ions are used up/ Reaction 2 is complete before the concentrations of the reactants changes (significantly)</p> <p>Allow so that the concentration of the reactants does not change(significantly)</p> <p>Ignore any references to the rate depending on the thiosulfate concentration</p> <p>Ignore references to the iodine/iodide being completely used up</p>	(1)

Question Number	Answer	Additional Guidance	Mark
2(a)(iv)	An answer that makes reference to the following point: <ul style="list-style-type: none"> so that the iodine disappears (in the reaction with the thiosulfate) before any reaction with the starch can occur 	<p>Allow arguments based on a slower rate for Reaction 2 e.g. the iodine would not be removed / the colour (of the complex) would appear too soon / straightaway/the colour change is delayed/not all the thiosulfate will react</p>	(1)

Question Number	Answer	Additional Guidance	Mark
2(b)(i)	<ul style="list-style-type: none"> concentration potassium iodide in mixture 	<p>Example of calculation</p> <p>$11 \times 0.200 \div 45 = 0.049 \text{ (mol dm}^{-3}\text{)}$</p> <p>Ignore SF except 1 SF</p>	(1)

Question Number	Answer	Additional Guidance	Mark
2(b)(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> axes, labels with units and scale chosen to cover at least half the graph in each direction 1st two and last two points correctly plotted (allow one small square) and best fit straight line. Ignore third point. 	<p>Example of correct graph</p> <p>Allow axes reversed</p>	(2)

Question Number	Answer	Additional Guidance	Mark
2(b)(iii)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> first order since the graph (of rate against concentration) is a straight line (through the origin)/ the rate/1/t is proportional to the concentration of iodide ions 	<p>M2 is dependent on M1 Allow the gradient is constant Ignore references to half-lives Do not award contradictory reasoning e.g. concentration is proportional to time</p>	(2)

Question Number	Answer	Additional Guidance	Mark
2(c)(i)	<p>An explanation that makes reference to two of the following points:</p> <ul style="list-style-type: none"> the temperature (1) the (total) volume of the reaction mixture (1) the volume of the KI (solution)/iodide ions (1) the volume of Na₂S₂O₃ (solution)/thiosulfate (1) 	<p>Accept just “total volume”</p> <p>Allow the volumes of other reactants for 1 mark Ignore references to concentration of individual reactants. Do not award the volume of peroxydisulfate Do not award pressure</p>	(2)

Question Number	Answer	Additional Guidance	Mark
2(c)(ii)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> rate = $k[\text{S}_2\text{O}_8^{2-}][\text{I}^-]$ 	<p>Allow $[\text{I}^-][\text{S}_2\text{O}_8^{2-}]$ Allow K</p>	(1)

Question Number	Answer	Additional Guidance	Mark
2(d)	<ul style="list-style-type: none"> • two simultaneous equations (1) • subtraction (1) • evaluation of activation energy with sign and units (1) <p>An alternative method:</p> <ul style="list-style-type: none"> • in a graph of $\ln k$ against $1/T$ the gradient gives $-E_a/R$ • gradient can be found by subtraction of $\ln k \div$ subtraction of $1/T$ • rearrangement to give E_a 	$-6.03 = - (E_a \times 0.00351 \div R) + \text{constant}$ $-3.47 = - (E_a \times 0.00314 \div R) + \text{constant}$ $-2.56 = - (E_a \times 0.00037 \div R)$ (+)57.496 kJ mol ⁻¹ / (+)57496/57500/57000 J mol ⁻¹ Ignore SF except 1 SF Correct answer with some evidence of working scores 3 Using values correctly rounded to 3SF (+)58181/58180/ 58200/58000 J mol ⁻¹ Using 1/285 and 1/318 gives (+)58425/58430/58400/58000 J mol ⁻¹ $\text{gradient} = \frac{-3.47 - (-6.03)}{0.00314 - 0.00351} = -6918.9 \text{ K}^{-1}$ $-(-6918.9 \times 8.31) = (+) 57.496 \text{ kJ mol}^{-1}$ Ignore SF except 1 SF	(3)

(Total for Question 2 = 15 marks)

Question Number	Answer	Additional Guidance	Mark
3(a)(i)	<p>A description that makes reference to three of the following points:</p> <ul style="list-style-type: none"> • add deionised/distilled water to the solid (in a beaker) (1) • (transfer solution to) a volumetric flask with washings (1) • make up to the mark (with deionised/distilled water) and invert/mix (1) 	<p>Allow the use of a funnel to transfer solid to flask</p> <p>Do not award conical flask/measuring cylinder</p> <p>M3 depends on the use of a volumetric flask in M2</p>	(3)

Question Number	Answer	Additional Guidance	Mark
3(a)(ii)	<p>An explanation that makes reference to two of the following points:</p> <ul style="list-style-type: none"> • volume measured using a measuring cylinder is less accurate/precise than a (volumetric) pipette (1) • the ethanedioic acid volume/amount is stoichiometric/used in the titration calculation/is a limiting factor (1) • the sulfuric acid acidifies the mixture/ provides hydrogen ions for the reaction/must be in excess (1) 	<p>Allow reverse argument</p> <p>Ignore references to a catalyst</p>	(2)










Question Number	Answer	Additional Guidance	Mark
3(a)(iii)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> the titration is self-indicating / the end-point is when the mixture becomes purple/pale pink 	<p>Allow colourless to pink Allow reactants and products are different colours Do not award pink to colourless</p>	(1)

Question Number	Answer	Additional Guidance	Mark
3(a)(iv)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> the reaction has a high activation energy / rate is slow (at room temperature) (1) the reaction will be faster at a higher temperature/heat is needed to supply the activation energy (1) Mn²⁺ ions are produced (in the titration reaction) (which) act as a catalyst/ the reaction is autocatalysed (1) 	<p>Allow the reaction is exothermic Allow Mn²⁺ ions speed up the reaction</p>	(3)


Question Number	Answer	Additional Guidance	Mark
3(a)(v)	<ul style="list-style-type: none"> • mol of manganate(VII) in titre • mol ethanedioic acid in 250 cm³ • mass water in solid • mole ratio of ethanedioic acid : water <p>Alternative M3: molar mass of ethanedioic acid</p> <p>Alternative M4: moles water</p>	<p>Example of calculation</p> <p>(1) $19.90 \times 0.0203 \div 1000 = 4.0397 \times 10^{-4} / 0.00040397$ (mol)</p> <p>(1) $0.00040397 \times 5/2 \times 10 = 1.00993 \times 10^{-2} / 0.010099$ (mol)</p> <p>(1) $1.27 - (0.010099 \times 90) = 0.36107$(g)</p> <p>(1) $0.0101 : 0.36107 \div 18$</p> <p>$0.0101 : 0.020059 / 2.0059 \times 10^{-2}$</p> <p>1: 2 so x=2</p> <p>Alternative M3 $1.27 \div 0.010099 = 125.75$</p> <p>$125.75 - 90 = 35.751$ $35.751 \div 18 = 1.9862 \sim 2$ so x=2</p> <p>Correct answer with some working scores 4 Ignore SF for M1,M2 and M3 Do not award M4 if answer is not an integer</p>	(4)

Question Number	Answer	Additional Guidance	Mark
3(b)	<p>An answer that makes reference to three of the following points:</p> <p>Method 1</p> <ul style="list-style-type: none"> • titrate (a solution of) ethanedioic acid with (aqueous) sodium hydroxide (noting the volume NaOH added at the end point) (1) • using thymol blue as an indicator (1) • add (25cm³) ethanedioic acid to the mixture and measure the pH using a pH meter (1) <p>Method 2</p> <ul style="list-style-type: none"> • add (small portions of) sodium hydroxide, noting the pH after each addition (1) • until the first neutralisation/ sharp rise in pH has been observed (1) • plot a graph of pH (against the volume of alkali added) and determine the point of half- neutralisation (1) 	<p>Ignore colour change even if incorrect</p> <p>Allow: add half the titre volume of NaOH to a different 25cm³ sample of ethanedioic acid, and measure the pH using a pH meter</p> <p>This could be determined by using any indicator but not phenolphthalein</p>	(3)

(Total for Question 3 = 16 marks)

Question Number	Answer	Additional Guidance	Mark						
4(a)	<p>An explanation that makes reference to three of the following points</p> <ul style="list-style-type: none"> corrosive toxic/acute toxicity/poisonous/fatal/lethal serious/long term health hazard/carcinogenic/cancer 	<table border="1"> <tr> <td></td> <td></td> <td></td> </tr> <tr> <td>corrosive</td> <td>toxic</td> <td>carcinogenic</td> </tr> </table> <p>2 correct scores 1 3 correct scores 2</p>				corrosive	toxic	carcinogenic	(2)
									
corrosive	toxic	carcinogenic							

Question Number	Answer	Additional Guidance	Mark
4(b)(i)	<p>An explanation that makes reference to two of the following points:</p> <ul style="list-style-type: none"> below 0°C the reaction is (too) slow above 5°C the diazonium salt/the product (of Reaction1) will decompose/HNO₂/nitrous acid is unstable 	<p>Accept phenol/nitrogen is formed Do not award the final product will decompose</p>	(2)

Question Number	Answer	Additional Guidance	Mark
4(b)(ii)	<p>A description that makes reference to two of the following points:</p> <ul style="list-style-type: none"> • (Buchner)funnel with perforated base and filter paper (1) • (Buchner) flask with side arm, seal(bung) and connection to (vacuum) pump/reduced pressure (1) <p>To score both marks both pieces of apparatus must be identifiable as separate</p>		(2)

Question Number	Answer	Additional Guidance	Mark
4(b)(iii)	<p>A description that makes reference to three of the following points:</p> <ul style="list-style-type: none"> • (place sample of solid in) a capillary/melting point tube (1) • insert tube into melting temperature apparatus Or Thiele tube/ oil bath with thermometer (1) • heat (to 110°C then slowly) till sample melts (1) 	<p>Accept a small tube/Pasteur pipette</p> <p>If a water bath is stated then M3 cannot be scored A labelled diagram may score M1 and M2 Ignore descriptions of recrystallisation</p>	(3)

Question Number	Answer	Additional Guidance	Mark
4(c)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> the solubility (in water) increases (so dye is lost from the body more easily/ the dye is more easily washed off the hands) 	Ignore any reasons given for the increased solubility even if incorrect.	(1)

(Total for Question 4 = 10 marks)

